Catalog No. MUIEF-01002-1

Lot Number: 130328C31



Certificate of Analysis

Strain ICR Mouse Embryonic Fibroblasts (Irradiated)

Cryopreservation Date: 2013-03-28

Passage Number: 1

Viability

Cells are assayed for viability post-thaw using vital staining assay with trypan blue.

Specification: Cells should exhibit $\geq 80\%$ viability.

Sterility

Bacterial and Fungal Contamination: Samples are inoculated and cultured on blood agar plate, thioglycolate broth, tryptocase soy broth and sabouraud dextrose agar.

Specification: No growth must be observed.

Mycoplasma: Samples are tested for mycoplasma contamination using a PCR-based assay and direct

culture.

Specification: Results must be negative.

Endotoxin: Samples are tested for endotoxin contamination with LAL test.

Specification: Results must show a concentration of ≤ 25 EU/ml.

Supporting Capability

Cells are shown to be able to support the growth of mouse embryonic stem (ES) cells and human ES and for the maintenance of the cells in undifferentiated state. The cells are used to culture mouse ESC and human ESC (in a separate set of plates) for three passages and each set of ESC are assayed for expression of cell-specific markers to confirm that cells remain undifferentiated. Immunostaining is performed using fluorochrome-conjugated antibodies specific to Oct4, SSEA-1, Nanog, SSEA-3, and SSEA-4.

Specification: For mouse ESC, the results indicate that $\geq 90\%$ of colonies in a plate, and > 90% of cells in each colony are positive for Oct4, SSEA-1, and Nanog, while $\leq 5\%$ of colonies in a plate, and < 5%of cells in each colony are positive for SSEA-3 and SSEA-4. For human ESC, the results must indicate that $\geq 90\%$ of colonies in a plate, and > 90% of cells in each colony are positive for Oct4, SSEA-3, SSEA-4, and Nanog, while $\leq 5\%$ of colonies in a plate, and < 5% of cells in each colony are positive for SSEA-1.



Verification of Growth Arrest

Cells are arrested with γ-ray. Cells are assayed after cryopreservation and confirmed to be mitotically inactivated. Cells are thawed and plated at low density (1.0–1.5 x 10⁴ cells/cm²) and monitored for cell growth for 14 days. After 14 days, the results indicate that no growth of cells is observed.

Results:

All specifications have been met.

Jane Chen QA Manager

Jane Chen

Apr 30, 2013