

Certificate of Analysis

Human Umbilical Cord Blood MSC

Catalog No. HUXUB-01001 Lot Number: 130606L01

Cryopreservation Date: 2013-06-06 Passage Number: 2

Viability

Cells are assayed for viability post-thaw using vital staining assay with trypan blue. Specification: Cells should exhibit $\geq 80\%$ viability.

Sterility

Bacterial and Fungal Contamination: Samples are inoculated and cultured in blood agar plate, thioglycolate broth, tryptocase soy broth and sabouraud dextrose agar. Specification: No growth must be observed.

Mycoplasma: Samples are tested for mycoplasma contamination using a PCR-based assay and direct culture.

Specification: Results must be negative.

Endotoxin: Samples are tested for endotoxin contamination with LAL test. Specification: Results must show a concentration of ≤ 10 EU/ml.

Exogenous Virus: Samples using ELISA assay to detect HIV, HBV, HCV and Syphilis. Specification: Results must be negative.

Purity

Cells are assayed for purity using flow cytometric analysis of cell surface antigen expression after cryopreservation. Cells are immunofluorescently stained with fluorochrome–conjugated antibodies specific to cell surface antigens CD29, CD105, CD44, CD45 and CD34. Specification: Cells must show \geq 70% positivity for expression of cell surface antigens CD29, CD44 and CD105. Cells must show \leq 5% positivity for expression of cell surface antigens CD34 and CD45.

Proliferation Ability

Cells are characterized by their ability to proliferate in culture with an attached well-spread morphology for ≥ 5 passages, and $\leq 5\%$ cells exhibit spontaneous differentiation in each passage.

Differentiation Ability

Cells are assayed after cryopreservation for their ability of tri-lineage differentiation. Cells must be able to differentiate to osteocytes, adipocytes and chondrocytes when cultured in the appropriate differentiation media.



Results:

All specifications have been met.

Jame chen

Jane Chen QA Manager Jul 27, 2013