

Certificate of Analysis

Strain C57BL/6 Mouse Embryonic Stem Cells

Catalog No. MUBES-01001

Lot Number: 140316L02

Cryopreservation Date: 2014-03-16

Passage Number: 20

Viability

Cells are assayed for viability post-thaw using vital staining assay with trypan blue.

Specification: Cells should exhibit $\geq 80\%$ viability.

Sterility

Bacterial and Fungal Contamination: Samples are inoculated and cultured on blood agar plate, thioglycolate broth, tryptocase soy broth and sabouraud dextrose agar.

Specification: No growth must be observed.

Mycoplasma: Samples are tested for mycoplasma contamination using a PCR-based assay and direct culture.

Specification: Results must be negative.

Endotoxin: Samples are tested for endotoxin contamination with LAL test.

Specification: Results must show a concentration of $\leq 25\text{EU/ml}$.

Karyotype

Cells are analyzed for karyotype by performing ≥ 20 metaphase spreads.

Specification: Results must be indicated that the cells possess 40 chromosomes (20 diploid pairs) plus 2 sex chromosomes (X, Y). The profile must match the published profile of Strain C57BL/6 mouse with no gross abnormalities.

Verification of Undifferentiated State

Cells are analyzed for expression of cell-specific markers after cryopreservation. Cells are immunostained with fluorochrome-conjugated antibodies specific to OCT-4, SSEA-1, Nanog, SSEA-3 and SSEA-4.

Specification: Cells must be shown to remain undifferentiated when cultured on mouse embryonic feeder cells after cryopreservation. Results must indicate that $\geq 90\%$ of colonies in a plate and $\geq 90\%$ of cells in each colony are positive for OCT-4, SSEA-1 and Nanog, while $\leq 5\%$ of colonies in a plate and $\leq 5\%$ of cells in each colony are positive for SSEA-3 and SSEA-4.

Differentiation Potential

Cells are assayed after cryopreservation for their ability to differentiate into embryoid bodies and express cell-specific markers indicative of the three germ layers.

Specification: Results must indicate positive expression of β 3-tubulin or nestin (ectoderm-specific markers), smooth muscle actin (mesoderm-specific marker), and AFP (endoderm-specific marker).

Results:

All specifications have been met.



Jane Chen
QA Manager
Jun 24, 2014